

-1-

DURATION OF MAGNETIC FIELD(S) ACTING ON METEORITE PARENT BODY(S).

M. W. Rowe
Department of Chemistry
Texas A&M University
College Station, Texas 77843

Study of remanent magnetization in meteorites began 15 years ago when Lovering and Stacey [1959] and Stacey and Lovering [1959] initiated their studies. Although attempts have been made at estimating the intensities of paleofields which acted on meteorite parent bodies [Stacey et al., 1961; Weaving, 1962; Gus'kova, 1963; Banerjee and Hargraves, 1972; Brecher, 1972; Butler, 1972], none have been directed toward estimating the temporal distribution or duration of these fields. It is the purpose of this note to undertake a preliminary attempt at estimating the time that the magnetic field(s) existed and acted on meteorite parent bodies. Information necessary for this preliminary analysis is available in the literature, although a more careful check on the assumptions used will require further studies--both on the diffusion of Ar^{40} and on the magnetic properties of meteorites.

For the present, I will confine my comments to magnetic studies on ordinary stony meteorites (chondrites) since recent studies indicate that paleointensity determinations on carbonaceous chondrites may be more complex than originally thought [Watson et al., 1974; Herndon et al., 1974] and necessary data is not available on the iron meteorites. There are eight chondrites which have had estimations of the intensity of the ancient fields implanting the magnetization in them: Brewster [Weaving, 1962], Farmington and Mt. Browne [Stacey et al., 1961], and Mordvinoka, Ochansk, Pultusk, Rakovka, and Zhovtnevyi [Gus'kova, 1963]. Of these, five have undergone the necessary chronological studies, i.e., gas retention ages: Farmington, Mt. Browne, Ochansk, Pultusk, and Zhovtnevyi [Zahringer, 1966]. K-Ar ages reported were 0.7, 4.1, 3.7, 3.9, and 4.0 Aeons ($\text{AE}=10^9$ years), respectively for those meteorites.

The essential argument is: If a sample exhibits a low K-Ar age, I assume that the sample was heated to a temperature high enough to drive off the requisite amount of Ar^{40} . By examining the rare gas studies arbitrarily selected from several laboratories on the thermal release of Ar^{40} from stone meteorites and lunar samples, I make a crude estimate of the temperature

(NASA-CR-138492) DURATION OF MAGNETIC
FIELD(S) ACTING ON METEORITE PARENT
BODY(S) (Texas A&M Univ.) 4 p HC \$4.00

CSCL 03B

G3/30

N74-26286

Unclas

40162

required to drive off a certain fraction of the Ar^{40} . After estimation of the temperature reached, I examine the effect of this temperature on the magnetization of the sample.

Of the five chondrites above which are dated, only Farmington is much lower than the expected age of ~4.6 AE with a K-Ar age of ~0.7 AE. Thus assuming a one-stage Ar^{40} loss I estimate that $\geq 87\%$ of the Ar^{40} must have been lost to result in such a low K-Ar age. To remove ~87% of the Ar^{40} from lunar samples by stepwise heating, temperatures of 1040 to 1100°C were required in three independent studies [Alexander *et al.*, 1972; Huneke *et al.*, 1972; Kaiser, 1972] and similar studies on four stone meteorites required temperature of 750 to 1200°C [Reynolds and Turner, 1964; Manuel *et al.*, 1972; Miller *et al.*, 1973]. The value of 750°C stands alone; all others were 1000°C or more. I thus estimate from this data that a temperature of ~1000°C is generally necessary to remove 87% of the Ar^{40} from diverse samples in laboratory studies. This is obviously a long extrapolation, i.e., from laboratory heating to parent body heating. But Wood [1967] presented good petrologic evidence that Farmington had been severely heated. Note, that the estimated temperature (~1000°C) is ~250°C higher than the Curie point of iron. While more experimentation is obviously desirable, new gas release studies via linear heating [Frick *et al.*, 1973] seem to support this as a reasonable means for estimating the temperature necessary for Ar^{40} removal. For example, Frick *et al.* found that if a sample is heated linearly to a certain temperature, T_m , allowed to cool and once again linearly heated to even higher temperatures, the gas release exhibits the following characteristic: Upon the second heating, only small amounts of additional Ar^{40} release are observed until about T_m is reached, where the Ar^{40} release increases rapidly until the rate of Ar^{40} release is virtually identical with the rate observed during the initial heating. Gas release at temperatures below T_m during the second heating is less than expected if the mechanism were purely classical diffusion loss.

Stacey, *et al.* [1961] demonstrated that the magnetization in Farmington was carried primarily by nickel-iron (as it was in all chondrites studied). Since the remanent magnetization in nickel-iron disappears at temperatures $\geq 770^\circ\text{C}$, no remanent magnetization is expected in Farmington since the

time of outgassing (~ 0.7 AE), unless it re-cooled in the presence of a magnetic field. Therefore I tentatively suggest that the Farmington chondrite parent body was outgassed ~ 0.7 AE ago (or more recently) losing $\geq 87\%$ of its total Ar^{40} and most likely reached at a temperature of $>770^\circ\text{C}$, thus erasing whatever magnetic remanence was previously recorded. However, a magnetic field seems to have been present on the Farmington parent body as recently as ~ 0.07 AE ago in view of the remanence found in the Farmington chondrite by Stacey, et al. Fig. 1 shows a schematic history of the Farmington meteorite.

That a magnetic field was associated with the Farmington meteorite as recently as ~ 0.7 AE ago should not be taken as evidence that a steady magnetic field existed in the early history of the solar system and extended to at least 0.7 AE ago. As Stacey [1967] aptly stated, "While alternative explanations of the primary magnetizations of chondrites cannot be finally excluded, the most satisfying ones at the present time require a parent body with a magnetic field. However, there is no experimental basis for asserting that it was a steady (terrestrial-type) field, generated by a well-developed core, and not a transient field accompanying the break-up of the body." Perhaps, in view of the difficulties involved in maintaining an ~ 4 AE long core on meteorite parent bodies, [Fish et al., 1960; Wood, 1967; Fricker et al., 1970; Herndon and Rowe, 1973], a transient field accompanying the break-up of the parent body is more probable than a steady field.

Acknowledgements. This work was supported by the National Science Foundation Grant No. GP-18716 and the National Aeronautics and Space Administration Grant No. NGR-44-001-152.

REFERENCES

- Alexander, Jr., E.C., P.K. Davis and R.S. Lewis, Science, **175**, 417-419, 1972.
Banerjee, S.K. and R.B. Hargraves, Earth Planet. Sci. Letters, **17**, 110-119, 1972.
Brecher, A., Proc. of Sym. "On the Origin of the Solar System", Nice, April 3-8, 260-273, 1972.
Butler, R.F., Earth Planet. Sci. Letters, **17**, 120-128, 1972.
Fish, R.A., G.G. Goles and E. Anders, Astrophys. J., **132**, 243-248, 1960.
Fricker, U., H. Baur, H. Funk, D. Phinney, C. Schafer, L. Schultz and P. Signer, Geochim. Cosmochim. Acta, Suppl. **4**, 1987-2002, 1973.
Fricker, P.E., J.I. Goldstein and A.L. Summers, Geochim. Cosmochim. Acta, **34**, 475-491, 1970.
Gus'kova, E.G., Geomagn. and Aeron., **3**, 308-312, 1963.

REFERENCES (Continued)

- Herndon, J.M. and M.W. Rowe, Nature (Phys. Sci.), 244, 40-41, 1973.
 Herndon, J.M., M.W. Rowe, E.E. Larson and D.E. Watson, to be submitted to Earth Planet. Sci. Letters, 1974.
 Huneke, J.C., F.A. Podosek, and G.J. Wasserburg, Earth Planet. Sci. Letters, 13, 375-383, 1972.
 Kaiser, W.A., Earth Planet. Sci. Letters, 13, 387-399, 1972.
 Lovering, J.F., Am. J. Sci., 257, 271-275, 1959.
 Manuel, O.K., R.J. Wright, D.K. Miller and P.K. Kuroda, Geochim. Cosmochim. Acta, 36, 961-983, 1972.
 Miller, D.K., D.W. Efurd, R.H. Thompson, M.A. Reynolds, K. Sakamoto and P.K. Kuroda, Personal communication, 1973.
 Reynolds, J.H. and G. Turner, J. Geophys. Res., 69, 3263-3281, 1964.
 Stacey, F.D., International Dictionary of Geophysics, Pergamon Press, N.Y., 1141-1143, 1967.
 Stacey, F.D. and J.F. Lovering, Nature, 183, 529-530, 1959.
 Stacey, F.D., J.F. Lovering and L.G. Parry, J. Geophys. Res., 66, 1523-1534, 1961.
 Watson, D.E., E.E. Larson, J.M. Herndon and M.W. Rowe, submitted to Earth Planet. Sci. Letters, 1974.
 Weaving, B., Geophys. J. Roy. Astron. Soc., 7, 203-211, 1962.
 Wood, J.A., Icarus, 6, 1-49, 1967.
 Zahringer, J., Meteoritika, 27, 25-40, 1966, in Russian.




Fig. 1. Schematic representation of the history of the Farmington meteorite. Time axis grossly distorted.